

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Chieftain® Duo 100EC

INSECTICIDE

ACTIVE CONSTITUENT: 100g/L ALPHA-CYPERMETHRIN
SOLVENT: 729 g/L LIQUID HYDROCARBONS

GROUP	3A	INSECTICIDE
-------	-----------	-------------



Control of insect pests of cereals, cotton, grain legumes, oilseeds, pastures, pome and stone fruits, trees & ornamentals, tobacco & vegetables as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 1 – 1000L

Sinon Australia Pty Limited
1.03/33 Lexington Drive
Bella Vista NSW 2153
ACN 101 741 024
TEL: 02 8883 0239

© Sinon registered Trademark

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray on any plants in flower while bees are foraging. This product is known to have a deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.

PROTECTION OF WILDLIFE , FISH , CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic invertebrates such as yabbies. DO NOT contaminate streams, rivers or waterways with the chemical or the used container. Drift and run-off from treated areas may be hazardous to fish or crustaceans in adjacent sites.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site.

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of water ways desirable vegetation and tree roots. Empty containers and product should not be burnt. If the container is a returnable/refillable type, when empty, or the contents are no longer required, return the container to the point of purchase. This type of container remains the responsibility and property of Sinon.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. Avoid inhaling vapour or spray mist. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and a face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126). If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

UN NO. 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS ALPHA- CYPERMETHRIN) MARINE POLLUTANT
PG III	HAZCHEM 2W



APVMA Approval No: **69532/61046**
Date of Manufacture:
Batch Number:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Chieftain Duo100EC
INSECTICIDE

ACTIVE CONSTITUENT: 100g/L ALPHA-CYPERMETHRIN
SOLVENT: 729 g/L LIQUID HYDROCARBONS

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

Control of insect pests of cereals, cotton, grain legumes, oilseeds, pastures, pome and stone fruits, trees & ornamentals, tobacco & vegetables as specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA APPROVAL NO: 69532/61046

Sinon Australia Pty Limited
1.03/33 Lexington Drive
Bella Vista NSW 2153
ACN 101 741 024
TEL: 02 8883 0239

© Sinon registered Trademark

DIRECTIONS FOR USE

Restrains: DO NOT APPLY IF RAINFALL IS EXPECTED WITHIN 6 HOURS OF APPLICATION

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS	
Cabbages Cauliflowers Broccoli, Brussels Sprouts, Kale, Kohl Rabi, Chinese Cabbage, Turnips	Cabbage white butterfly (<i>Pieris rapae</i>), Cabbage moth (<i>Plutella xylostella</i>) <i>Helicoverpa spp.</i>	All states	Low volume 400 mL/ha	1 day (harvest)	Apply when pest populations indicate. When reinfestation is continuous, treatment every 7-10 days may be required. Add a non-ionic surfactant at its label rates. LOW VOLUME: Ground rig application: Apply in 100 to 600L water per hectare as a fine spray with droplet size of 100 to 200 microns For aerial application, apply in 20 to 60 L water/ha with a droplet size of 100 to 150 microns HIGH VOLUME use a spray with a droplet size of 200 to 400 microns. Apply 600 L spray mixture per hectare just after transplanting and increase gradually to 1000L/ha toward maturity. ULTRA LOW VOLUME: see ULV application section of this label. <i>Helicoverpa armigera</i> In Qld and NSW – follow the application directions for the pest above. Apply as required for pest incidence. Thorough and frequent crop checks are essential. Preferably apply to eggs. Apply to larvae only if they are less than 5mm long.	
	Cluster caterpillar (<i>Spodoptera litura</i>)	NSW, Vic, WA & ACT	High Volume 50 mL/100L ULTRA LOW VOLUME: 400mL/ha			
Canola	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA, Tas & WA only	200 mL/ha or 300mL/ha	21 days (cutting for harvest or stock feed or grazing)	Do not use more than a total of 400mL/ha per season to any one crop. For ULTRA LOW VOLUME use, see application section of this label. Inspect the crop regularly and immediately after flowering. Apply when damaging pest numbers first appear on the crop and repeat if necessary. For aerial application, use a total volume of 30-35 L/ha and apply in cooler part of the day. Use the higher rate on larvae larger than 10mm. Crops should be inspected as they merge. Border sprays are required to control invading adults. Apply when cotyledons and leaves are being eaten or the plant topped. Repeat as necessary. Apply according to pest incidence.	
	Tobacco lopper (<i>Chrysodexis argentifera</i>)	NSW, Tas, SA & Vic only				
	Vegetable weevil (<i>Listroderes difficilis</i>)	NSW, ACT, Vic, Tas, SA, WA only	400 mL/ha			
	Cabbage white butterfly (<i>Pieris rapae</i>), cabbage moth (<i>Plutella xylostella</i>)					
	Red legged earth mite (<i>Halotydeus destructor</i>)	All States except NT and Qld	100mL/ha			Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor red legged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.
	Red legged earth mite (<i>Halotydeus destructor</i>), blue oat mite (<i>Penthaleus major</i>)		50mL/ha			Apply when mite numbers reach damaging levels. Do NOT apply as pre-emergence treatment. Do not use as a ULV application.

Cotton	Native budworm (<i>Helicoverpa punctigera</i>),	Qld, NSW, NT & WA only	300mL/ha	14 days (harvest)	For Ultra Low Volume use see ULV application section of this label. Apply as indicated by field checks using rates appropriate for the infestation level determined. Application should be timed to coincide with egg hatching and before larvae are in protected feeding sites.
			400mL/ha		Apply when there are up to 75 eggs and/or up to 5 larvae less than 5mm long per 100 terminals.
			500mL/ha		Apply when there are up to 150 eggs and/or up to 10 larvae less than 5mm long per 100 terminals and/or when larvae between 5 and 10mm are present.
	Cotton bollworm (<i>Helicoverpa armigera</i>)	Qld, NSW, NT & WA only	300mL/ha		Apply when there are up to 150 eggs and/or more than 10 larvae less than 5mm long per 100 terminals and/or when larvae longer than 10mm are present.
			400mL/ha		Preferably apply to eggs. Apply to larvae only if they are less than 5mm long.
			500mL/ha		Apply when there are up to 75 eggs and/or up to 5 larvae less than 5mm long per 100 terminals.
	Rough bollworm (<i>Earias huegeli</i>)	NSW, NT, Qld, WA only	300 or 400 mL/ha		Apply when there are up to 150 eggs and/or up to 10 larvae less than 5mm long per 100 terminals.
					Apply when there are more than 150 eggs and/or more than 10 larvae less than 5mm long per 100 terminals.
					Apply when 2 or more larvae are present per 100 bolls. It is essential to detect and treat infestations in the early stages before larvae are established or concealed in bolls deep in the canopy. Use the higher rate if larvae greater than 10 mm are present. Best results are obtained by applying at egg hatch.
Field Peas	Native budworm (<i>H. punctigera</i>)	WA only	160mL/ha	4 weeks (harvest)	For Ultra Low Volume use, see ULV application section of this label. Apply to open, less dense crops when damaging numbers of newly hatched larvae first appear on the crop and repeat if necessary.
		NSW, Tas, Vic, Act, SA & WA only	200 or 300mL/ha		Apply when damaging pest numbers first appear on the crop and repeat if necessary. Use the higher rate if larvae longer than 10mm are present. Best results are obtained by applying at egg hatch.

Field peas (cont)	Pea weevil (<i>Bruchus pisorum</i>)	NSW, ACT, Vic, SA & WA only	160 or 200mL/ha	4 weeks (harvest)	Apply during flowering prior to egg laying when adult weevil population reaches 1 or more per 25 sweeps of a sweep net. Use the higher rate for longer residual protection.
	Cutworm (<i>Agrostis</i> spp.)		75 mL/ha		Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Apply product in late afternoon or evening.
	Red legged earth mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA & WA only	100mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor red legged earth mite populations and re-treat as necessary. DO NOT use as a ULV application.
	Red legged earth mite (<i>Halotydeus destructor</i>), blue oat mite (<i>Penthaleus major</i>)		50mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as pre-emergence treatment. Do not use as a ULV application.
Grapevines (non bearing)	Pink cutworm (<i>Agrostis munda</i>), apple weevil (Curculio beetle) (<i>Otiorynchus cribricollis</i>), garden weevil (<i>Phlyctinus callosus</i>)	NSW, SA, Vic, Tas, WA only	100mL/100L	-	Monitor young vines during spring and early summer and apply at the first signs of leaf damage. Spray the leaves, canes and soil around each vine to a diameter of 30cm. 70 to 80mL of dilute spray should be sufficient for each vine. If pest infestation persists, a second application may be required after 3 weeks.
Lettuce	<i>Helicoverpa</i> spp.	All States	Low volume: 400mL/ha High volume: 50mL/100L	3 days (harvest)	Thoroughly and regularly check the crop. Apply at the first sign of pest activity. Preferably apply to eggs. Apply to <i>H. armigera</i> ONLY if larvae are less than 5mm long. Repeat according to pest incidence.
Linola	Native budworm (<i>H. punctigera</i>)	WA only	160 or 200mL/ha	12 weeks (harvest)	DO NOT apply more than a total 400mL/ha per season to any one crop. For Ultra Low Volume use, see ULV application section of this label. Inspect crop regularly during and immediately after flowering. Apply when damaging pest numbers first appear on the crop. For aerial application, apply during the cooler part of the day in a total volume of 30-35L/ha. Use the higher rate if larvae longer than 10mm are present. Refer to application section for water rates.

Linseed	Cutworms (<i>Agrostis</i> spp.)	NSW, ACT, SA, Tas only	75 mL/ha	14 days (harvest)	Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Apply product in late afternoon or evening.
	Native budworm (<i>H. punctigera</i>)	NSW, Vic, Tas, SA, WA only	200 or 300mL/ha		For Ultra Low Volume use, see ULV application section of this label. Inspect the crop regularly and immediately after flowering. Apply when damaging numbers first appear on the crop and repeat if necessary. Use the higher rate if larvae larger than 10 mm are present. Best results will be obtained by spraying at egg hatch. Refer to application section for water rates.
Lucerne (seed and forage crops)	Native budworm (<i>H. punctigera</i>)	NSW, Vic, Tas, SA, WA only	160mL/ha	14 days (grazing or cutting for stock feed)	For Ultra Low Volume use, see ULV application section of this label. Do NOT apply more than one application per cut or grazing for animal feed. Apply when pest populations reach economically damaging levels. Apply to larvae less than 5mm in length.
	Green mirid (<i>Creontiades dilutis</i>)				Do NOT apply more than one application per cut or grazing for animal feed. Apply when pest populations reach economically damaging levels.
Lupins	Native budworm (<i>H. punctigera</i>)	NSW, ACT, Vic & SA only	200 or 300mL/ha	4 Weeks (harvest)	DO NOT apply more than a total 600mL/ha per season to any one lupin crop. For Ultra Low Volume use, see ULV application section of this label. Apply when damaging pest numbers first appear on the crop and repeat if necessary. Use the higher rate if larvae larger than 10 mm are present. Best results are obtained by spraying at egg hatch.
		WA only	120 or 200mL/ha		Spraying should be timed to precede the first visible damage to the pods. Use the higher rate when the infestation is severe, or when residual activity is required.
	Cutworm (<i>Agrostis</i> spp.)	NSW, ACT, Vic, Tas, SA, WA only	75 mL/ha		Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Apply product in late afternoon or evening.
	Common armyworm (<i>Mythimna convecta</i>), Southern armyworm (<i>Persectania ewingii</i>)	ACT, NSW only	240mL/ha		Spray in the cool of the day (late afternoon) when larvae are most active.

Lupins (cont.)	Red legged earth mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	100mL/ha	4 Weeks (harvest)	Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor red legged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.
	Red legged earth mite (<i>Halotydeus destructor</i>), blue oat mite (<i>Penthaleus major</i>)		50mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as pre-emergence treatment. Do not use as a ULV application.
Maize	Corn earworm (<i>H. armigera</i>)	Qld, NSW ACT, Vic, NT only	300 or 400mL/ha	7 days (harvest)	For Ultra Low Volume use, see ULV application section of this label. Thoroughly and regularly check the crop. Apply from early silking according to pest incidence. Use the higher rate if larvae longer than 10mm are present. In Qld, NSW & NT, preferably apply to eggs or apply to larvae only if they are less than 5 mm long.
	Native budworm (<i>H. punctigera</i>)	All States	300 or 400mL/ha	7 days (harvest)	Thoroughly and regularly check the crop. Apply when the infestation reaches economically damaging levels and repeat as required. Best results will be obtained by applying at egg hatch. Use the higher rate if larvae longer than 10mm are present.
Beans (Mung and Navy)	Native budworm (<i>H. punctigera</i>)	Qld, NSW, ACT & NT only	300 or 400mL/ha	7 days (harvest)	For Ultra Low Volume use, see ULV application section of this label. Thoroughly and regularly check the crop. Small larvae are easier to kill than large larvae. Apply when the number of larvae feeding on flowers or pods reaches 1 to 2 per metre of row. Repeat as required. Best results will be obtained by applying at egg hatch. Use the higher rate if larvae longer than 10mm are present or when canopy is dense.
	Corn earworm (<i>H. armigera</i>)				Thoroughly and regularly check the crop. Apply when the infestations reach economically damaging levels and repeat as required. Preferably apply to eggs. In NSW & Qld, apply to larvae only if they are less than 5mm long. Use the higher rate when pest pressure is high.

Pastures (legume and grass based pastures)	Wingless grasshoppers (<i>Phaulacridium vittatum</i>)	All States	160mL/ha	3 days (grazing) 14 days (cut for stockfeed)	Do NOT apply more than a total of 320mL/ha per season. For Ultra Low Volume use, see ULV application section of this label. Apply to infested areas and repeat as necessary. Spraying is most effective on newly emerged hoppers before they begin dispersing. Spray in the warmer parts of the day when hoppers are exposed. Later sprays should be applied before the start of egg laying. Good coverage is essential.
	Brown pasture looper (<i>Ciampa arietaria</i>)	NSW, Vic, Tas, SA & WA only	50mL/ha		Apply when pest population reaches a commercially damaging level.
	Blackheaded pasture cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA only	100mL/ha		Spraying is most effective when larvae are detected and treated early. Suspect paddocks should be dug after the first substantial rain in April/May and inspected to ensure grubs are present in sufficient numbers to warrant treatment. Spraying after June will give poorer results.
	Red legged earth mite (<i>Halotydeus destructor</i>)	NSW, ACT, Vic, Tas, SA, WA only	100mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor red legged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.
	Red legged earth mite (<i>Halotydeus destructor</i>), blue oat mite (<i>Penthaleus major</i>)		50mL/ha		Apply when mite numbers reach damaging levels. Do not use as a ULV application. Autumn/winter: Apply 4 to 7 weeks after the opening rains in late autumn/ early winter when RLEM are present (2-3 weeks after egg hatch occurs). This product is rainfast after spray deposits have dried on the leaf surface. This product can be mixed with herbicides used for winter cleaning of sub-clover pastures. See the "compatibility" section of this label. Spring: If RLEM/BOM numbers increase in the spring, spray when damage is observed and again before diapause egg production begins. This product can be mixed with herbicides used for spray topping pastures. See the "compatibility" section of this label. Do NOT use as a pre-emergence treatment.
Pome fruit: Apples, pears	Apple weevil (<i>Otiorynchus cribricollis</i>) Garden weevil (<i>Phlyctinus callosus</i>)	NSW, Vic, SA, WA only	100mL/100L water	14 days (harvest)	Spray approximately 1-2 litres of solution onto the crotch, trunk and the soil at the base of each tree at peak weevil emergence. This usually occurs in late October – late November for garden weevil, and late November – mid December for apple weevil. Monitor weevil emergence using a single sided cardboard trunk band. Continue monitoring after spraying as a second spray may be needed 3-4 weeks later.

Rice (both aerial and drill sown)	Common armyworm (<i>Mythimna convecta</i>)	NSW only	200mL/ha	7 days	Do NOT apply more than a total of 400mL/ha per season to any one crop. Inspect crops regularly for the presence of grubs from flowering onwards. Apply when rice-damaging pest numbers first appear. Apply by aircraft in 20-30 litres of water per hectare to drained fields only. Spray in the cool of the day (early morning or late afternoon) when larvae are most active. Monitor crops closely and re-treat if necessary. Poor control may occur in crops that have lodged. See application section for correct water rates.
Soybeans	Native budworm (<i>H. punctigera</i>)	Qld, NSW, WA, ACT & NT only	300 or 400mL/ha	7 days	For Ultra Low Volume use, see ULV application section of this label. Thoroughly and regularly check the crop. Apply when flower or pod feeding numbers reach 1-2 per metre of row. Repeat as required. Use the higher rate if larvae longer than 10mm are present. Best results are obtained by applying at egg hatch.
	Corn earworm (<i>H. armigera</i>)				Thoroughly and regularly check the crop. Apply when the numbers are sufficient to cause economic damage. Preferably apply to eggs. In NSW and Qld, apply to larvae only if they are less than 5mm long. Repeat as required. Use the higher rate when pest pressure is high.
Stone fruit: Apricots, nectarines, peaches, plums	Apple weevil (<i>Otiorynchus cribricollis</i>) Garden weevil (<i>Phlyctinus callosus</i>)	WA only	100mL/100L water	14 days (harvest)	Spray approximately 1-2 litres of solution onto the crotch, trunk and the soil at the base of each tree at peak weevil emergence. This usually occurs in late October – late November for garden weevil, and late November – mid December for apple weevil. Monitor weevil emergence using a single sided cardboard trunk band. Continue monitoring after spraying as a second spray may be needed 3-4 weeks later.
Sorghum	Native budworm (<i>H. punctigera</i>), Corn earworm (<i>H. armigera</i>)	Qld, NSW, ACT & NT only	300 or 400 L/ha	7 days (harvest)	For Ultra Low Volume use, see ULV application section of this label. Crop checking should commence when the head emerges from the boot and continue at daily intervals until the end of flowering for midge and at weekly intervals until maturity for <i>H. armigera</i> . DO NOT apply to tight headed varieties. Apply when there are 2 or more actively feeding larvae per head, or when numbers are sufficient to cause economic damage. Use the higher rate if longer residual control is required. Preferably apply to eggs. Apply to <i>H. armigera</i> larvae only if they are less than 5mm long. Repeat as required.
	Sorghum midge (<i>Contarinia sorghicola</i>)		100 or 200mL/ha		Apply when midge numbers reach 1 -2 per head, from head emergence to completion of flowering. Repeat as required. Use the higher rate for longer residual protection.

Sunflowers	Native budworm (<i>H. punctigera</i>),	Qld, NSW, Vic, ACT & NT only	300 or 400mL/ha	21 days (harvest)	TO PROTECT BEES and ensure adequate pollination, application during flowering should be avoided. If application is necessary at flowering, apply early morning or late afternoon when bees are not actively foraging. For Ultra Low Volume use, see ULV application section of this label. Crop checking should be aimed to detect larvae as they hatch. Small larvae are easier to kill than large larvae. Apply when an average of 2-3 larvae are present per head or when economic damage is occurring. Repeat as required. Apply before the heads turn downwards to ensure adequate coverage. Use the higher rate when larvae larger than 10 mm are present. Best results will be obtained by applying at egg hatch.
	Corn earworm (<i>H. armigera</i>)				Thoroughly and regularly check the crop. Apply when numbers are sufficient to cause economic damage. Preferably apply to eggs. In NSW and Qld apply to larvae only if they are less than 5 mm long, repeat as required. Use the higher rate under heavy pest pressure.
	Grey cluster bug (<i>Nysius clevelandensis</i>), Rutherglen bug (<i>Nysius vinitor</i>)				Apply from budding when adult numbers per plant reach 10-15 in dryland crops and 20-25 in irrigated crops. After flowering apply when adult numbers on the face of heads reaches 20-25. Repeat as required. The higher rate should be used when numbers are very high.
	Rutherglen bug (<i>Nysius vinitor</i>)	Vic, Tas, WA only	250mL/ha	Apply from budding when adult numbers per plant reach 10-15 in dryland crops and 20-25 in irrigated crops. After flowering apply when adult numbers on the face of heads each 20 -25. Repeat as required.	
Sweet corn	Native budworm <i>H. punctigera</i>), Corn earworm (<i>H. armigera</i>)	All States	300 or 400mL/ha	7 days (harvest)	For Ultra Low Volume use, see ULV application section of this label. Thoroughly and regularly check the crop. Cob damage tolerated is variable according to market requirements. For fresh market corn spray at tassel emergence then at intervals of 5-8 days until silks wither. For processing corn and maize apply at early silking. Larvae in protected feeding sites within the cob are not effectively controlled. Apply before this situation occurs. Best results will be obtained by applying at egg hatch. Use the higher rate if larvae larger than 10 mm are present. To help contain pyrethroid resistance in <i>Helicoverpa armigera</i> in summer crops, do NOT apply to corn earworm longer than 5mm.

Tobacco	Native budworm (<i>Helicoverpa punctigera</i>) Tobacco budworm (<i>H. armigera</i>)	Vic only	30 or 40mL/100L	7 days (harvest)	Apply from just after transplanting on a 7 to 10 day schedule, according to pest incidence. Apply as a medium to fine spray using hollow and/or solid cone nozzles. The spray volume should be gradually increased as the plants grow, from 200L/ha just after transplanting to 1000L/ha at maturity. Use the higher rate when larvae longer than 10mm are present or when egg laying is intense.	
Tomatoes (bush and trellis)	Native budworm (<i>H. punctigera</i>)	All States	UTLRA LOW VOLUME: 300 OR 400mL/ha	1 day (harvest)	<p>Do NOT apply to trellis tomatoes by aircraft. Apply on a 7 day schedule while pests are active. Use the middle rate when pest activity is high and/or larvae between 10 and 20mm in length are present. Use the highest rate when larvae longer than 20mm are present and/or when interruption of the schedule enables a very severe infestation to develop.</p> <p>ULTRA LOW VOLUME: see ULV application section of this label. LOW VOLUME: By ground rig: apply in 100 to 400L of water per hectare as a fine spray. By aircraft: apply in a minimum of 10L of water per hectare as a fine spray of 100 to 150 microns VMD. HIGH VOLUME: Apply as a medium to fine spray. Gradually increase the spray volume as the plants grow, from 200L/ha just after transplanting establishment to 1000L/ha at maturity.</p>	
	Tomato grub (<i>H. armigera</i>)	Vic, Tas, SA, WA only	LOW VOLUME: 200, 300 or 400mL/ha			
	Cluster caterpillar (<i>Spodoptera litura</i>)	Qld, NSW, ACT, WA, NT only	HIGH VOLUME: 20,30 or 50mL/100L			
	Tomato grub (<i>H. armigera</i>)	Qld, NT, NSW only	UTLRA LOW VOLUME: 300 mL/ha LOW VOLUME: 300mL/ha HIGH VOLUME: 30mL/100L			Thoroughly check the crop the crop at 2-3 day intervals from transplanting/emergence. Apply according to incidence. Preferably apply to eggs. Apply to larvae only if they are less than 5mm long. Apply using the methods described for native budworm above.
	Plague thrips (<i>Thrips imaginis</i>)	Qld, NSW, ACT, Vic, Tas, WA, NT only	UTLRA LOW VOLUME: 130mL/ha LOW VOLUME: 130mL/ha HIGH VOLUME: 18mL/100L			The crop should be frequently checked when it is flowering for the presence of the pest. Apply when the infestation reaches an economically damaging level using the application methods as described above.

Winter cereals	Cutworms (<i>Agrotis spp.</i>)	NSW, ACT, Vic, SA, WA only	75mL/ha	7 days (harvest) 14 days (stubble grazing)	Do NOT apply more than a total of 540 mL/ha per season to any one crop. For Ultra Low Volume use, see ULV application section of this label. Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Spray in the late afternoon or evening.
		Qld only	75 or 150mL/ha		In Qld use the higher rate when the infestation is severe, or when the larvae are larger than 10 mm long, or when the residual activity is required.
	Pasture Webworm (<i>Hednota spp.</i>)	NSW, Vic, SA, WA only	75mL/ha		Do NOT use as a ULV application. Pre-planting: May be applied pre-planting with knockdown herbicides. Apply from the last week of May when larvae have emerged. Do NOT apply to dense pasture. All pasture should be closely grazed prior to application to ensure adequate spray penetration. Apply in a minimum of 100L of water per hectare. Apply at first sign of pest infestation. Repeat as necessary. Post Crop Emergence: Inspect crop regularly from emergence and apply at the first sign of pest activity. Repeat as required.
	Red legged earth mite (<i>Halotydeus destructor</i>)	NSW, ACT, Vic, Tas, SA, WA only	100mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor red legged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.
	Red legged earth mite (<i>Halotydeus destructor</i>), blue oat mite (<i>Penthaleus major</i>)		50mL/ha		Apply when mite numbers reach damaging levels. Spray seedling crops if silvering or whitening (bleaching) of the leaves is causing a reduction in crop growth. If possible spray on a calm, mild morning when mites are actively feeding on crop leaves. Do NOT apply as pre-emergence treatment. Do not use as a ULV application.
	Aphids (<i>Rhopalosiphum spp.</i>) (barley yellow dwarf virus vectors)		125mL/ha		To control aphids, sprays should be applied at 3 and 7 weeks after emergence to reduce aphid colonization and spread of Barley Yellow Dwarf virus. This will also reduce the effect of feeding aphid damage.
	Common armyworm (<i>Mythimna convecta</i>), Southern armyworm (<i>Persectania ewingii</i>)	All States	240mL/ha		Apply before "head lopping" occurs when larval numbers exceed 2 or more per square metre. Spray in the cool of the day (late afternoon) when the larvae are most active. Spray to achieve good crop penetration. This rate is effective against larvae up to 20 mm in length. Monitor crops regularly and re-treat if necessary. Poor control may occur in crops that have lodged. See application section for correct water rates.

Trees & Ornamentals

Eucalyptus	Adults and larvae of eUcalyptus leaf beetle (<i>Chrysophtharta bimaculata</i>)	Tas only	250 mL/ha	-	Apply by fixed wing aircraft or by helicopter using hydraulic nozzles or micronair equipment, to the crowns of eucalypt trees. Micronair application in 5 litres of water/ha of water has proved effective. Apply before insect damage causes severe defoliation. Treatment will control small and large larvae as well as adult beetles. For ULV application, see ULV application section of this label.
Banksias Ornamentals	Banksia moth (<i>Danima banksiae</i>)	WA only	20 mL/100L	-	Apply on a regular program at 2 week intervals at early flower development. Commence spraying when blooms are immature and continue until flowers are fully developed.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

BROCCOLI, BRUSSELS SPROUTS, CABBAGES, CAULIFLOWERS, CHINESE CABBAGE, KALE, KOHLRABI, TOMATOES, TURNIPS - DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

LETTUCE - DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

MAIZE, MUNG BEANS, NAVY BEANS, RICE, SORGHUM, SOYBEANS, SWEET CORN, TOBACCO: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

**WINTER CEREALS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
DO NOT GRAZE TREATED STUBBLE FOR 14 DAYS AFTER APPLICATION**

LUCERNE - DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

COTTON, LINSEED, STONE & POME FRUIT: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

**CANOLA: DO NOT CUT AND WINDROW FOR HARVEST FOR 21 DAYS AFTER APPLICATION
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 21 DAYS AFTER APPLICATION.**

SUNFLOWERS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION

FIELD PEAS & LUPINS: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION

**LINOLA: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION
PASTURES - DO NOT GRAZE FOR 3 DAYS AFTER APPLICATION.
- DO NOT CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

This product is a contact and residual insecticide. It can be used as a protective treatment when applied at regular intervals or as a knockdown treatment to control existing larvae. Best results will be obtained by spraying at egg hatch. Thorough coverage is essential to ensure adequate control. Apply during the cooler parts of the day.

The product can be applied mixed with water carrier OR oil based bulking agents such as D-C-Tron Cotton Spray Oil or other compatible ULV products.

MIXING

Low Volume and High volume applications by ground rig or aircraft when the product is applied with water as a carrier

Add the required quantity to water in the spray tank and mix thoroughly. Maintain agitation during mixing and application.

Ultra Low Volume (ULV) application by aircraft when the product is applied with oil based bulking agents
This product can be mixed with D-C-Tron Cotton Spray Oil or other compatible products (see Compatibility Section).

Add the mixing partner to the spray tank first, engage agitation system and add the required amount of Chieftain Duo 100EC Insecticide direct to the spray tank. DO NOT mix with water to ensure there is no water in the spraying system.

APPLICATION

Low volume and High Volume by ground rig or aircraft when Chieftain Duo 100EC Insecticide is applied with water carrier

This product can be applied by ground or aircraft with a water carrier. Thorough coverage is essential to ensure adequate control. Always apply with a non-ionic surfactant at its labeled rate, unless detailed on the label of the tank mix partner. Apply in cooler parts of the day or night.

Ground Application – water carrier : For low volume spraying of field crops with ground rigs, use a total volume of 50-200 L/ha except for sweet corn, tomatoes and tobacco where higher volumes should be used. Drop arms should be used on ground rigs in row crops taller than 300mm. The application should be made as a fine spray, preferably using hollow cone nozzles and a droplet size of 150-200 microns unless otherwise recommended in the Critical Comments.

Aerial Application – water carrier :

DO NOT apply to trellis tomatoes by aircraft.

Use at least 20 L/ha of total spray volume.

For spring/early summer applications to cereals, linola, canola, rice and to other dense crops, apply in a total spray volume of 30 to 35L/ha. If possible, spray in a cross wind. Avoid spraying in calm conditions or when wind is light and variable in direction. Apply as a spray of 100-150 microns VMD.

Ultra Low Volume (ULV) application by aircraft

Chieftain Duo 100EC Insecticide mixed with D-C-Tron Cotton Spray Oil or other compatible products should be applied in a minimum total spray volume of 1.5L/ha. It should only be applied by aircraft with suitable equipment to provide a droplet size of approximately 80-100 microns VMD. Applications should be made during the cooler parts of the day or at night. Avoid application in calm or very windy conditions. Preferably apply in light to moderate cross winds.

COMPATIBILITY

Low Volume and High Volume Application by ground rig or aircraft when Chieftain Duo 100EC Insecticide is applied with water as a carrier

This product is compatible with D-C-Tron Cotton Spray Oil, Mancozeb 750 WG, dicamba, Kocide¹, methomyl, chlorpyrifos 300, Ridomil¹, Wuxal¹, Clethodim, , dimethoate, diquat, glyphosate, Tigrex¹, Jaguar¹, simazine, Imazathyr 700 WG, 2,4-D Amine, 2,4-D Ester, 2,4-DB, MCPA, paraquat.

Do NOT mix this product with wettable powders and water dispersible granules BEFORE addition to the spray tank.

This product can be mixed with Mancozeb 750 WDG providing the mixture is agitated efficiently and used immediately.

Ultra Low Volume by aircraft

This product should only be mixed with specific ULV formulations of other insecticides eg Chlorpyrifos 300, and PBO synergists, when mixed according to the directions on the PBO synergist labels.

INSECTICIDE RESISTANCE WARNING

GROUP	3A	INSECTICIDE
-------	----	-------------

For insecticide resistance management Chieftain Duo 100EC Insecticide is a Group 3A insecticide.

Some naturally-occurring insect biotypes resistant to Chieftain Duo 100EC Insecticide and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Chieftain Duo 100 EC Insecticide or other Group 3A Insecticides are used repeatedly. The effectiveness of Chieftain Duo 100EC Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Sinon Australia Pty Limited accepts no liability for any losses that may result from the failure of Chieftain Duo 100EC Insecticide to control resistant insects.

Chieftain Duo 100EC Insecticide may be subject to specific resistance strategies. For further information contact your local supplier, Sinon representative or local department of agriculture agronomist.

In NSW and Qld, application of this product to *Helicoverpa armigera* larvae longer than 5mm may not only be ineffective but it may increase the level of synthetic pyrethroid resistance. This product should NOT be used to treat infestations that were not controlled by an earlier application of it or another synthetic pyrethroid. Infestations not controlled by this product should be treated with another insecticide from another chemical group. Application of this product with an insecticide from another chemical group such as mehtomyl will assist with the management of synthetic pyrethroid resistant *Helicoverpa armigera*.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray on any plants in flower while bees are foraging. This product is known to have a deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.

PROTECTION OF WILDLIFE , FISH , CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic invertebrates such as yabbies. DO NOT contaminate streams, rivers or waterways with the chemical or the used container. Drift and run-off from treated areas may be hazardous to fish or crustaceans in adjacent sites.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site.

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of water ways desirable vegetation and tree roots. Empty containers and product should not be burnt. If the container is a returnable/refillable type, when empty, or the contents are no longer required, return the container to the point of purchase. This type of container remains the responsibility and property of Sinon.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. Avoid inhaling vapour or spray mist.

When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and a face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126). If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of

merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

¹ Not trademark of Sinon