

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

Sinon Chloro 900 WG Fungicide

ACTIVE CONSTITUENT: 900 g/kg CHLOROTHALONIL

GROUP	M5	FUNGICIDE
--------------	-----------	------------------

For the Control of Fungal Diseases on Almonds, Apricots, Bananas, Carrots, Celery, Cherries, Chickpeas, Faba Beans, Grapes, Lentils, Onions, Peaches, Peanuts, Peas, Plums, Potatoes, Tomatoes and Vegetables as per the Directions for Use

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 1 kg to 20 kg

Sinon Australia Pty Ltd
ACN 102 741 024
Level 7, 60 York Street, Sydney NSW 2000
TEL: 03 5441 8907

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool-well ventilated area. DO NOT store for prolonged periods in direct sunlight

Plastic containers: Plastic containers: Triple-rinse containers before disposal. Add rinsings to the spray tank.

DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will damage eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. If product in eyes, wash out immediately with water. If clothing becomes contaminated with product or wet with spray, remove clothing immediately. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26; New Zealand 0800 764 766.* If swallowed, DO NOT induce vomiting.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

APVMA Approval No: 90436/131215

BN:

DOM:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

Sinon Chloro 900 WG Fungicide

ACTIVE CONSTITUENT: 900 g/kg CHLOROTHALONIL

GROUP	M5	FUNGICIDE
--------------	-----------	------------------

For the Control of Fungal Diseases on Almonds, Apricots, Bananas, Carrots, Celery, Cherries, Chickpeas, Faba Beans, Grapes, Lentils, Onions, Peaches, Peanuts, Peas, Plums, Potatoes, Tomatoes and Vegetables as per the Directions for Use

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No: 90436/131215

Sinon Australia Pty Ltd
ACN 102 741 024
Level 7, 60 York Street, Sydney NSW 2000
TEL: 03 5441 8907

DIRECTIONS FOR USE

Restraints

DO NOT mix spraying oils or wetting agents with this product.

For all stone fruit except cherries, the following additional restraints apply:

DO NOT use oils after bud-swell and during the season if Sinon Chloro 900 WG Fungicide is to be used after shuck fall.

DO NOT tank mix Sinon Chloro 900 WG Fungicide with EC formulations when spraying after shuck fall.

DO NOT exceed 2.5 kg of Sinon Chloro 900 WG Fungicide per hectare of crop.

1) TREE AND VINE CROPS

RATE In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.					Critical Comments For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Situation & Crop	Disease	States	Rate	WHP (days)	
Almonds	Shot-hole (<i>Stigmina carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)	SA, Vic, NSW, WA & Tas only	130 g/ 100 L Apply 2000 L/ha	-	Apply at bud-swell, bud-burst, pink bud, shuck fall, cap fall. Apply every 10 - 14 days. Apply 1 week pre-harvest.
Apricots	Brown rot- Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)			7	Apply at bud-swell, bud-burst, pink bud, full bloom. Apply no later than 35 days pre-harvest. See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmina carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>) Freckle (<i>Venturia carpophila</i>)			Apply at bud-swell, bud-burst, pink bud, shuck fall, cap fall, then every 10 - 14 days. Spray applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits	
Cherries	Brown rot - Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)	Qld, SA, Vic, NSW, WA & Tas only	130 g/ 100 L Apply 2000 L/ha	7	Apply at bud-swell, bud burst, pink bud, early blossom, full bloom. Apply 3 weeks pre-harvest and 1 week pre-harvest.
	Shot-hole (<i>Stigmina carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)	All states		Apply at bud-swell, bud burst, pink bud, shuck fall, cap fall, then every 10 - 14 days. Apply 1 week pre-harvest.	
	Transit rot (<i>Rhizopus stolonifer</i>)			Apply 1 week pre-harvest	

RATE In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.					Critical Comments For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Situation & Crop	Disease	States	Rate	WHP (days)	
Grapes Note: Russeting of some table grape varieties may occur	Downy mildew (<i>Plasmopara viticola</i>) Bunch rot (<i>Botrytis cinerea</i>)	All states	1.5 – 1.9 kg/ha	Dessert 7 Wine 14	Spray at first appearance of the foliage disease. Apply at 7 - 14 day intervals particularly during wet weather and periods of rapid growth. For bunch rot, applications at early full bloom, two weeks later, at veraisson and two weeks prior to harvest. Table grapes: Russeting of some varieties may occur. DO NOT apply after cap fall on these varieties.
	Black Spot (Anthracnose) (<i>Eisinoe ampelina</i>)		175 g/ 100 L		
Nectarines	Shot-hole (<i>Stigmina carpophila</i>)	All states except Qld	133 g/ 100 L Apply 2000 L/ha	7	Apply at bud-swell, pink bud, shuck fall and then repeat each 14 days if weather conditions favour disease. Apply no later than 35 days pre-harvest as unacceptable fruit skin damage may occur. See also Additional Restraints for stone fruits.
	Brown rot- Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)		Apply at bud-swell, pink bud, early blossom and full bloom. Apply no later than 35 days pre-harvest as unacceptable fruit skin damage may occur. See also Additional Restraints for stone fruits.		

RATE In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.					Critical Comments For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Situation & Crop	Disease	States	Rate	WHP (days)	
Peaches	Brown rot- Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)	All states	130 g/ 100 L Apply 2000 L/ha	7	Apply at bud-swell, pink bud, early blossom and full bloom. Apply no later than 35 days pre-harvest as unacceptable fruit skin damage may occur See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmia carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)				Apply at bud-swell and continue at 7 - 14 day intervals. Apply no later than 35 days pre-harvest as unacceptable fruit skin damage may occur. See also Additional Restraints for stone fruits.
	Leaf curl (<i>Taphrina deformans</i>)				Apply at bud-swell and continue at 7 - 14 day intervals. In Qld spray only at bud-swell. Apply no later than 35 days pre-harvest as unacceptable fruit skin damage may occur See also Additional Restraints for stone fruits.
Plums	Brown rot- Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)	Qld, SA, Vic, NSW, WA & Tas only	133 g/ 100 L Apply 2000 L/ha	1	Apply at bud-swell, bud-burst, early blossom, pink bud and full bloom. Spray applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmia carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)	All states			Apply at bud-swell, bud burst, shuck fall and cap fall. Spray application later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.

2) LEGUMES

Situation & Crop	Disease	State	Rate	WHP (days)	Critical Comments
Beans- Broad (Faba)	Chocolate spot (<i>Botrytis fabae</i>) Rust (<i>Uromyces viciae-fabae</i>)	All states	1.2 - 1.9 kg/ha	7	Apply at first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 7 - 14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.
Chickpeas Lentils	Ascochyta blight (<i>Aschochyta rabeie</i>) Ascochyta blight (<i>Aschochyta lentis</i>), Grey Mould (<i>Botrytis fabae</i>)	Qld, NSW, Vic, SA, WA only	0.8-1.6 kg/ha	14 (harvest and grazing)	<p>Preplant treatment of Chickpeas or Lentils seed with P-Pickel T will control transmission of Ascochyta and Botrytis. It is therefore advised as part of an overall program to manage these diseases in Chickpeas and Lentils. Foliar applications of Sinon Chloro 900WG as indicated below will provide ongoing control of these diseases in these crops. Apply in a minimum total water volume of 100L/ha for ground application and 300L/ha for aerial application.</p> <p>Use the lower application rate in the following circumstances:</p> <ol style="list-style-type: none"> At the early stage of crop growth when there is less vegetation to cover and band spraying is possible. In low disease pressure situations where there is no history of Ascochyta infection in the paddock or adjacent paddocks and/or seasonal conditions are dry. Where a strict 2 week interval is practiced in higher disease pressure situations. <p>Use the higher application rate in the following circumstances:</p> <ol style="list-style-type: none"> In high disease pressure where there is an existing infection or infected stubble is present in the paddock or adjacent paddocks and regular rainfall events are occurring. On highly susceptible varieties, especially where the application interval has extended beyond 2 weeks <p>Commence application at 3 to 5 weeks after emergence in Chickpeas and 8 to 10 weeks after emergence in Lentils and make following applications at 2 to 4 week intervals depending on disease presence or prior to rainfall events.</p> <p>Use the shorter spray intervals when:</p> <ol style="list-style-type: none"> In high disease pressure situations (infection present and regular rainfall events) or When higher susceptible varieties are being grown. <p>The longer spray intervals may be used when:</p> <ol style="list-style-type: none"> No rain has occurred since the last application or When the higher application rate has been used in the previous application and where the variety has low to moderate susceptibility to Ascochyta or Grey Mould infection.

Situation & Crop	Disease	State	Rate	WHP (days)	Critical Comments
					<p>If applying prior to rainfall events:</p> <ol style="list-style-type: none"> 1. Reapply if 2 or more weeks have elapsed since the last application. 2. Apply Hemani Chlorothalonil 900WG at 0.8 kg/ha if less than 4 weeks have elapsed since the last application and the variety being grown. 3. Apply Hemani Chlorothalonil 900WG at 1.6 kg/ha if 4 or more weeks have elapsed since the last application and a highly susceptible variety is being grown or infection is present in the crop. <p>If Grey Mould is the predominant disease in Lentils and infection pressure is high on a susceptible variety, tank mix or alternate Hemani Chlorothalonil 900WG WITH A Botrytis specific fungicide such as carbendazim , at critical times such as mid-flowering to mid-pod fill.</p>
Peanuts	<p>Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>) Peppery leaf spot (<i>Leptosphaerulina trifolii</i>) Peanut rust (<i>Puccinia arachidis</i>) Net blotch (<i>Didymosphaeria arachidicola</i>)</p>	Qld, NSW, WA & NT only	900 g – 1.5 kg/ha	-	<p>Use higher rate when conditions favour leaf spot or when rust appears. DO NOT feed peanut vines to livestock. NSW AND WA ONLY: Commence application when disease first appears and repeat as necessary to maintain control.</p> <p>Qld ONLY: NORTH AND SOUTH BURNETT: Commence spraying when disease appears and repeat when rain and warm moist nights are expected. ATHERTON TABLELANDS: Commence spraying 4-6 weeks after planting and repeat at 7 - 14 day intervals depending on conditions.</p>
Peas	Downy mildew (<i>Peronospora viciae</i>)	Qld, WA & Tas only	900 g – 1.5 kg/ha	7	<p>Apply at first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 7 -14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.</p>

3) VEGETABLES AND FRUIT

Situation & Crop	Disease	State	Rate	WHP (days)	Critical Comments
Artichoke-Globe	Grey Mould (<i>Botrytis cinerea</i>)	All states	1.5 - 1.9 kg/ha	1	Apply at 7 - 10 day intervals as required. Use the higher rate if disease pressure is high.
Bananas DO NOT apply to bananas within 14 days of applying an oil based spray or crop damage will occur	Leaf Spot (<i>Mycosphaerella musicola</i>)	Qld, NSW, WA, NT only	1.5 kg/ha	1	For North Qld: Spray every 14 days late January to end of March. Extend to 21 days after March and then 28 - 35 days in the drier months of the year. For South Qld: Spray every 21 days from start of December to end of April. For NSW: Spray every 21 days from end of November to end of May. For WA: Spray every 10 - 14 days from November to end of May and extend to 14 - 21 days from May to November.
		WA only	1.3 kg/ha		
	Leaf speckle (<i>Mycosphaerella musae</i>)	Qld, NSW, WA, NT only	2.5 kg/ha		
Broccoli, Brussels sprouts, Cabbages, Cauliflowers	Ring Spot (<i>Mycosphaerella brassicicola</i>)	All states	1.5 – 3.0 kg/ha	7	Apply at first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 14 day intervals. Use higher rate if disease pressure is high.
Capsicums Peppers	Grey mould (<i>Botrytis cinerea</i>) Anthracnose (<i>Colletotrichum ascutatum</i>)		1.5 - 1.9 kg/ha	1	Apply at 7 - 10 day intervals as required. Use the higher rate if disease pressure is high
Carrots	Leaf Blight (<i>Alternaria dauci</i>)		1.5 kg/ha	7	Commence application as soon as symptoms favour the development of the disease and then spray every 7-14 days.
Celery	Septoria leaf spot (<i>Septoria apiicola</i>)		1.5 kg/ha	1	Apply every 7 - 14 days. Apply also to seed beds
	Cercospora early blight (<i>Cercospora apii</i>) (NSW, WA only)	NSW & WA only	1.0 kg/ha		Apply every 10 - 14 days
Cucurbits (including Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Rockmelon,	Downy Mildew (<i>Pseudoperonospora cubensis</i>) Gummy stem blight (<i>Didymella bryoniae</i>) Anthracnose (<i>Coffetotrichum orbiculare</i>) Leaf blight (<i>Alternaria cucumerina</i>) Target leaf spot (<i>Cercospora citruffina</i>)	All states	1.3 – 2.1 kg/ha	1	Apply on a 7 to 14 day schedule. Apply rates towards the higher end of the recommended range at 7 day intervals when disease pressure is high. Commence spray program early, at first true leaf and before vining or flowering. When used at these rates, this product will suppress powdery mildew.

Situation & Crop	Disease	State	Rate	WHP (days)	Critical Comments
Watermelon, Zucchini)	Belly rot (<i>Rhizoctonia solani</i>)		1.3 – 2.1 kg/ha		Suppression of belly rot will be obtained where fruit form on soil when sprayed using the above foliar spray program.
Endive	Grey mould (<i>Botrytis cinerea</i>)	All states	1.5 – 1.9 kg/ha	1	Apply at 7 - 10 day intervals as required. Use the higher rate if disease pressure is high
Leeks, Shallots	Purple blotch (<i>Alternaria porri</i>)		1.5 – 1.9 kg/ha	1	Apply at first appearance of disease. Repeat at 7 - 10 day intervals, particularly during wet weather.
Okra	Leaf diseases/spot	Qld & WA only	1.5 – 1.9 kg/ha	1	Apply at first appearance of disease. Repeat at 7 - 10 day intervals.
Onions (excluding spring onions)	Downy mildew (<i>Peronospora destructor</i>)	All states	1.9 kg/ha	14	Apply at first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 14 day intervals.
Potato	Early blight or Target spot (<i>Alternaria solani</i>) Late blight or Irish blight (<i>Phytophthora infestans</i>)		900 g – 1.5 kg/ha	-	Spray at first sign of foliage disease or at flowering time, whichever is earlier. Spray every 7 - 14 days until within 14 days of desiccation or harvest.
Radish	Grey mould (<i>Botrytis cinerea</i>)		1.5 – 1.9 kg/ha	1	Apply at 7 - 10 day intervals as required. Use the higher rate if disease pressure is high.
Rhubarb	Ascochyta leaf spot (<i>Ascochyta rhei</i>)		120 g/ 100 L Apply 2000 L/ha	7	Apply every 10 – 14 days.
Sweet corn	Turcicum leaf blight (<i>Exserohilum turcicum</i>)		1.5 – 1.9 kg/ha	1	Apply at first appearance of disease. Repeat at 7 - 10 day intervals.
Tomato	Early blight or Target blight (<i>Alternaria solani</i>) Leaf mould (<i>Fulvia fulva</i>) Grey leaf spot (<i>Stemphylium solani</i>) Botrytis rot or Grey mould (<i>Botrytis cinerea</i>)		1.5 – 1.9 kg/ha	1	Spray at first appearance of the foliage disease. Spray at 7 - 14 day intervals. Continue into the picking season to obtain best results against leaf mould and Botrytis rot.
Watercress	Cercospora leaf spot		Qld & WA only	1.5 – 1.9 kg/ha	1

4) MISCELLANEOUS

Situation & Crop	Disease	State	Rate	WHP (days)	Critical Comments
Ornamentals Chrysanthemums, Geraniums, Iris & other ornamentals	Botrytis, Grey mould, Rust diseases, Septoria and Alternaria leaf spot diseases	All states	80 g/ 100 L	-	Apply at 7 - 14 day intervals. Use 7-day interval under severe conditions. Avoid application to blooms in azalea, hydrangea and petunia.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

HARVEST	
Artichoke, Bananas, Capsicum, Celery, Cucurbits, Endive, Leek, Okra, Peppers, Plums, Radish, Shallots, Sweet Corn, Tomatoes, Watercress	DO NOT HARVEST FOR 1 DAY AFTER APPLICATION
Apricots, Beans, Broccoli, Brussels Sprouts, Cabbages, Carrots, Cauliflowers, Cherries, Nectarines, Peaches, Peas, Rhubarb, Table Grapes	DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
Chickpeas, lentils, onions, wine grapes	DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION
Almonds, Potatoes, Peanuts	NOT REQUIRED WHEN USED AS DIRECTED
GRAZING	
Chickpeas, Lentils	DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION
Beans, Peas, Peanuts	DO NOT GRAZE LIVESTOCK ON TREATED CROPS
DO NOT GRAZE OR FEED TREATED FORAGE OR FODDER TO LIVESTOCK WHICH MAY BE USED TO PRODUCE MILK FOR HUMAN CONSUMPTION	

GENERAL INSTRUCTIONS

Sinon Chloro 900 WG Fungicide is a protectant fungicide for the control of a range of diseases in fruit and vegetables. For effective disease control a good leaf cover must be achieved before a widespread disease outbreak occurs.

APPLICATION

Dilute Spraying:

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying:

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only:

1. Dilute spray volume as determined above: For example 1500 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 3 x (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)
4. If the dilute label rate is 1 kg/ 100 L, then the concentrate rate becomes 3 x 1, that is 3 kg/ 100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For further information on concentrate spraying, users are advised to consult the relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

MIXING

Half fill spray vat and with agitation system running, add the required amount of product. Top up with water to the required volume. The product must be kept in suspension at all times by continuous agitation. Where prepared spray mixes have been allowed to stand, thoroughly re-agitate before using.

Application rates per hectare:

Low Volume - 200 litres or more.

High Volume - 500 litres or more.

Aerial Application - 20 litres or more.

Mister - 75 to 255 litres.

The volume of water applied should increase as the plants grow in size to maintain adequate cover.

COMPATIBILITY

This product is compatible with wettable powder and flowable formulations of most commonly used fungicides, insecticides and miticides. Unless experience in your area has shown other combinations to be physically compatible and non-injurious to your crop do not combine with other chemicals, especially oil-based emulsifiable or flowable pesticides. This product should not be mixed with spraying oils or sprayed onto crops that have been sprayed with oil for at least 10 days after the last spray.

Wetting Agents: The addition of surfactants does not improve this product's efficiency and under some conditions may cause plant injury.

FUNGICIDE RESISTANCE WARNING

GROUP	M5	FUNGICIDE
--------------	-----------	------------------

Sinon Chloro 900 WG Fungicide is a member of the multi-site activity group of fungicides. For fungicide resistance management the product is a Group M5 fungicide. Some naturally occurring individual fungi resistant to Sinon Chloro 900 WG Fungicide and other Group M5 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Sinon Chloro 900 WG Fungicide or other Group M5 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Sinon Australia Pty Ltd accepts no liability for any losses that may result from the failure of Sinon Chloro 900 WG Fungicide to control resistant fungi.

PRECAUTION

Enclosed cabs (with closed windows and filters) for airblast application. DO NOT use human flaggers unless they are protected by engineering controls such as enclosed cabs.

Re-entry - DO NOT re-enter treated areas until the spray has dried. If prior entry is required, wear chemical resistant gloves.

PROTECTION OF LIVESTOCK

DO NOT contaminate *feed* or foodstuffs with this product or empty containers.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool-well ventilated area. DO NOT store for prolonged periods in direct sunlight

Plastic containers: Plastic containers: Triple-rinse containers before disposal. Add rinsings to the spray tank.

DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will damage eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. If product in eyes, wash out immediately with water. If clothing becomes contaminated with product or wet with spray, remove clothing immediately. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26; New Zealand 0800 764 766.* If swallowed, DO NOT induce vomiting.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.