



Product Name: ERASER 750 INSECTICIDE
APVMA Approval No: 59572/126195

Label Name:	ERASER 750 INSECTICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	750 g/kg ACEPHATE
Mode of Action:	GROUP 1B INSECTICIDE
Statement of Claims:	For the control of certain insect pests on Bananas, Crucifers, Macadamias, Ornamentals, Potatoes and Tomatoes as per the Directions for Use table.
Net Contents:	1kg - 20kg 3kg (6 x 500 g) Contains 6 x 500 g measure packs which it is illegal to sell separately 5kg (10 x 500 g) Contains 10 x 500 g measure packs which it is illegal to sell separately 3kg (1 x 3kg measure pack)
Restraints:	
Directions for Use:	This section contains file attachment.

Other Limitations:	
--------------------	--

Withholding Periods:	WITHHOLDING PERIOD: BRUSSELS SPROUT, CABBAGE, CAULIFLOWERS, POTATOES AND TOMATOES – DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. BROCCOLI – DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION. BANANAS, MACADAMIAS, ORNAMENTALS – NOT REQUIRED WHEN USED AS DIRECTED.
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	GENERAL INSTRUCTIONS MIXING Add the required amount of Eraser 750 Insecticide to the full volume of water in the spray tank with agitator in operation. Ensure the powder is completely dispersed before spraying. NOTE: A varying degree of consolidation of the contents of this container may occur during transport but the net weight of the contents is not changed and is as stated on the label. APPLICATION METHOD By ground: Crops should be sprayed with sufficient volume to ensure adequate coverage of plants. For high volume spraying, apply in 500 to 1000 litres water per hectare. By air: Apply in 10 to 45 litres of water per hectare. COMPATIBILITY Eraser 750 Insecticide may be mixed in the spray vat with any of the following products: Copper Oxychloride, Zineb. A mixture of Eraser 750 Insecticide with other products may be ineffective or may cause serious damage and hence is NOT recommended.
-----------------------	---

Resistance Warning:	INSECTICIDE RESISTANCE WARNING GROUP 1B INSECTICIDE For insecticide resistance management Eraser 750 Insecticide is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Eraser 750 Insecticide and other Group 1B Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. The effectiveness of Eraser 750 Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Sinon Australia Pty Limited accepts no liability for any losses that may result from the failure of Eraser 750 Insecticide to control resistant insects.
---------------------	--

Precautions:	
--------------	--

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS Eraser 750 Insecticide MAY CAUSE FOLIAGE DAMAGE TO CERTAIN SHRUBS AND TREES IE Crab Apple, Cotton wood, Lombardy poplar and American elm. DO NOT spray more than once every 28 days on carnations and chrysanthemums. Certain
--------------	---

	chrysanthemum varieties ie Showoff, Iceberg, Albatross, Bonnie Jean and Statesman have, on occasions, been damaged by products containing acephate. PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT spray while bees are actively foraging on and around the treatment area. Avoid spray drift from target area. DO NOT contaminate streams, rivers or waterways with this product or the used containers.
Storage and Disposal:	Store in the original container, tightly closed in a safe well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Bag in a Box - Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product. Water soluble bags in box - Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.
Safety Directions:	Product and spray is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.
First Aid Instructions:	If swallowed, splashed on skin or in eyes, or inhaled, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.
First Aid Warnings:	

DIRECTIONS FOR USE

CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			per 100L water	per ha		
Bananas (Bell Injection and bunch spray only)	Banana Scab moth (<i>Nacoleia octasema</i>), Banana flower thrips (<i>Thrips hawaiiensis</i>)	Qld, NSW, WA, NT only	168g (17g/10L)	-	-	Apply 30mL of prepared solution to each emerging banana bell. Inject the required volume directly into the upright bell as it emerges from the throat of the banana plant.
	Banana rust thrips (<i>Chaetanapho- thrips signipennis</i>)					Apply to the point of run-off, up to 50mL of prepared spray solution to each bunch, ensuring thorough, even spray coverage. Application should be made no later than 2 weeks after bunch emergence.
Brussels Sprouts	Heliothis (<i>Helicoverpa</i> spp.) including corn earworm and native budworm	Tas, NT only	130g	1.3kg	3 days (broccoli 14 days)	Spray when infestation occurs and then at 10 to 14 day intervals or as necessary. Add wetting agent to solution at rate of 10ml per 100 litres.
	Cabbage moth, cabbage white butterfly	NSW, ACT, Vic, SA, Tas WA, NT only	130g	1.3kg		
	Corn earworm, native budworm	Qld, WA only	100g	1kg		
	Cabbage aphid	Qld, WA, NT only	100g	1kg		
Cabbages, cauliflowers, broccoli	Cabbage moth, cabbage white butterfly	NSW, ACT, Vic, Tas, SA, WA, NT only	130g	1.3kg		
	Cabbage aphid, corn earworm, native budworm	Qld, WA, NT only	100g	1kg		
Macadamias	Flower eating caterpillar, fruitspotting bug, macadamia leaf miner, banana spotting bug, flower thrips	Qld, NT only	66g	660g	-	Apply at early flowering if pests are evident. One to three sprays may be required depending on the time and severity of the infestation. Spray when bees are least active.
	Monolepta beetle	Qld, WA, NT only	100g	1kg		Apply when activity first observed and repeat at 14 to 21 day intervals or as necessary.
	Fruitspotting bug, macadamia flower caterpillar, macadamia leaf miner, Macadamia nutborer	NSW, WA only				

Ornamentals	Aphids	All States	130g	1.3kg	-	Spray when infestation occurs and then at 14 day intervals. DO NOT apply to carnations later than 6 days before picking.
	Corn earworm, Native budworm	Vic, SA, Tas, WA only				
Potatoes	Potato moth (tobacco leafminer)	All States	66g	660g	3 days	Spray every 14 to 21 days or as necessary.
Tomatoes	Aphids	NSW, ACT, Vic, SA, WA, Tas only	100g	1kg	3 days	Spray when infestation occurs and then at 7 to 14 day intervals or as necessary (ground application only)
	Potato moth, Heliothis (<i>Helicoverpa</i> spp.)	NSW, ACT, Vic, SA, WA, Tas only	130g	1.3kg		
	Potato moth, Heliothis (<i>Helicoverpa</i> spp.), aphids	Qld, NT only	100g	1kg		Apply when infestation occurs then at 7 day intervals or as necessary.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**